arXiv:1502.05302v2 [math.AP] 17 May 2016

A Proof of Schiffer's Conjecture in Starlike Domain by Far-Field Patterns

Lung-Hui Chen[1]

May 18, 2016

Abstract

We formulate the Schiffer's conjecture in spectral geometry in the context of scattering theory. The problem is equivalent to finding a non-trivial solution in an interior transmission problem. We compare the back-scattering data of the perturbation along all incident angles. The uniqueness of the inverse scattering problem along each incident direction proves the Schiffer's conjecture.

MSC: 35P25/35R30/34B24.
Keywords: inverse scattering/Helmholtz equation/Rellich's lemma/interior transmission eigenvalue /Cartwright-Levinson theory.

1 Introduction

In this paper, we study the following inverse spectral problem:

$$\begin{cases} \Delta u + k^2 u = 0, & \text{in } D,\, k^2 \in \mathbb{R}^+; \\ \frac{\partial u}{\partial \nu} = 0, & \text{on } \partial D; \\ u = 1, & \text{on } \partial D, \end{cases} \tag{1.1}$$

where ν is the unit outer normal; D is a fixed starlike domain in $\mathbb{R}^3$ containing the origin with Lipschitz boundary ∂D. We interpret the model as the plane waves perturbed by the boundary condition which is specified by D, and satisfies the Helmholtz equation outside D. Let u be a non-trivial eigenfunction with some $k^2 \in \mathbb{R}^+$. We want to show that D are actually balls centered at origin.

Here we prove the result as a special case of interior transmission problem [2, 4, 10, 11, 12, 13, 14, 15, 16, 17, 18, 21, 22, 24, 25, 28, 29, 31]. In interior transmission problems, we look for a frequency so that a perturbed stationary wave behaves like or somewhere like a spherical Bessel function outside the perturbation. In Schiffer's conjecture, we ask if there is a frequency so that a perturbed wave can stay in its initial shape traveling to infinity in constant speed. We refer to [1, 28] and the reference there for the connections of interior transmission problem to other questions in mathematical science.

To give a point of view from scattering theory to (1.1), we take the incident wave field to be the time harmonic acoustic plane wave of the form

$$u^i(x) := e^{ikx\cdot d},$$

$k \in \mathbb{R}^+$, $x \in \mathbb{R}^3$, and $d \in \mathbb{S}^2$ is the incident direction. The inhomogeneity is defined by the index of refraction $n \in \mathcal{C}^2(\mathbb{R}^3)$ of (1.1), and the wave propagation is governed by the following equation.

$$\begin{cases} \Delta u(x) + k^2 n(x) u(x) = 0,\ x \in \mathbb{R}^3; \\ u(x) = u^i(x) + u^s(x),\ x \in \mathbb{R}^3 \setminus D; \\ \lim_{|x|\to\infty} |x|\{\frac{\partial u^s(x)}{\partial |x|} - iku^s(x)\} = 0, \end{cases} \tag{1.2}$$

---

[1] Department of Mathematics, National Chung Cheng University, 168 University Rd. Min-Hsiung, Chia-Yi County 621, Taiwan. Email: mr.lunghuichen@gmail.com; lhchen@math.ccu.edu.tw. Fax: 886-5-2720497.

in which the third equation is the Sommerfeld's radiation condition. Particularly, we have the following asymptotic expansion on the scattered wave field [17, 20].

$$u^s(x) = \frac{e^{ik|x|}}{|x|} u_\infty(\hat{x}; d, k) + O(\frac{1}{|x|^{\frac{3}{2}}}),\ |x| \to \infty, \tag{1.3}$$

which holds uniformly for all $\hat{x} := \frac{x}{|x|}$, $x \in \mathbb{R}^3$, and $u_\infty(\hat{x}; d, k)$ is known as the scattering amplitude or far-field pattern in the literature [17, 21]. It has an expansion in spherical harmonics [17, p. 35, Theorem 2.15]

$$u_\infty(\hat{x}; d, k) = \frac{1}{k} \sum_{n=0}^{\infty} \frac{1}{i^{n+1}} \sum_{m=-n}^{n} a_n^m Y_n^m(\hat{x}), \tag{1.4}$$

where we follow the notation in the reference.

Let us start with the Rellich's representation in scattering theory. We expand the possible solution u of (1.1) in a series of spherical harmonics near infinity by Rellich's lemma [17, p. 32, p. 227]:

$$u(x; k) = \sum_{l=0}^{\infty} \sum_{m=-l}^{m=l} a_{l,m}(r) Y_l^m(\hat{x}), \tag{1.5}$$

where $r := |x|$, $r \geq R_0$ with a sufficiently large R_0; $\hat{x} = (\theta, \varphi) \in \mathbb{S}^2$. The summations converge uniformly and absolutely on suitable compact subsets away from D. The spherical harmonics

$$Y_l^m(\theta, \varphi) := \sqrt{\frac{2l+1}{4\pi} \frac{(l-|m|)!}{(l+|m|)!}} P_l^{|m|}(\cos\theta) e^{im\varphi},\ m = -l, \ldots, l;\ l = 0, 1, 2, \ldots, \tag{1.6}$$

form a complete orthonormal system in $\mathcal{L}^2(\mathbb{S}^2)$, in which

$$P_n^m(t) := (1-t^2)^{m/2} \frac{d^m P_n(t)}{dt^m},\ m = 0, 1, \ldots, n, \tag{1.7}$$

where the Legendre polynomials P_n, $n = 0, 1, \ldots$, form a complete orthogonal system in $L^2[-1, 1]$. We refer this to [17, p. 25]. By the orthogonality of the spherical harmonics, the family of functions

$$\{u_{l,m}(x; k)\}_{l,m} := \{a_{l,m}(r) Y_l^m(\hat{x})\}_{l,m} \tag{1.8}$$

satisfy the first equation in (1.1) independently for each (l, m) in $r \geq R_0$ for sufficiently large R_0.

Now we consider the boundary condition given by the second and third equations in (1.1), and then extend the solutions $u_{l,m}(x; k)$ into $r \leq R_0$ as follows. Let $\hat{x}_0 \in \mathbb{S}^2$ be any given incident direction that intersects ∂D at $(\hat{R}, \hat{x}_0) \in \mathbb{R}^+ \times \mathbb{S}^2$. For the given $\hat{x}_0$, we impose the differential operator

$$\Delta = \frac{1}{r^2} \frac{\partial}{\partial r} r^2 \frac{\partial}{\partial r} + \frac{1}{r^2 \sin\varphi} \frac{\partial}{\partial \varphi} \sin\varphi \frac{\partial}{\partial \varphi} + \frac{1}{r^2 \sin^2\varphi} \frac{\partial^2}{\partial \theta^2}$$

on $u_{l,m}(x; k)$ and, accordingly, we have the following ODE:

$$\frac{d^2 a_{l,m}(r)}{dr^2} + \frac{2}{r} \frac{d a_{l,m}(r)}{dr} + (k^2 - \frac{l(l+1)}{r^2}) a_{l,m}(r) = 0, \tag{1.9}$$

which is solved by spherical Bessel functions and spherical Neumann functions. Let

$$y_{l,m}(r) := r a_{l,m}(r), \tag{1.10}$$

so we obtain

$$\begin{cases} y''_{l,m}(r) + (k^2 - \frac{l(l+1)}{r^2}) y_{l,m}(r) = 0; \\ y_{l,m}(0) = 0. \end{cases} \tag{1.11}$$

To give an initial condition, we apply the boundary conditions in (1.1) to $u_{l,m}(x;k)$ near the intersection points $\hat{R}$ along $\hat{x}_0$. We replace the boundary condition $\frac{\partial u}{\partial \nu}=0$ to be $\nabla u=0$. Hence,

$$[\frac{y_{l,m}(r;k)}{r}]'|_{r=\hat{R}}=0; \tag{1.12}$$

$$[\frac{y_{l,m}(r;k)}{r}]|_{r=\hat{R}}=1, \tag{1.13}$$

in which we assume there is no tangent point. The solutions $y_{l,m}(r;k)$ are independent of m, so we write $y_{l,m}(r;k)$ as $y_l(r;k)$. Hence, now we have following boundary conditions.

$$\frac{y_l'(\hat{R};k)}{\hat{R}}-\frac{y_l(\hat{R};k)}{\hat{R}^2}=0; \tag{1.14}$$

$$y_l(\hat{R};k)-\hat{R}=0. \tag{1.15}$$

If k satisfies (1.14) and (1.15), then $y_l'(\hat{R};k)=1$. Thus, (1.12) and (1.13) equivalently satisfy

$$F_l(k;\hat{R}):=y_l(\hat{R};k)-\hat{R}=0; \tag{1.16}$$

$$G_l(k;\hat{R}):=y_l'(\hat{R};k)-1=0. \tag{1.17}$$

In the initial state, $y_l(\hat{R};k)$ is exactly the boundary defining function of D. Combining (1.11), (1.16), and (1.17), we consider the following eigenvalue problem at $\hat{R}$ for each fixed $\hat{x}\in\mathbb{S}^2$ and all $l\geq 0$:

$$\begin{cases} y_l''(r;k)+(k^2-\frac{l(l+1)}{r^2})y_l(r;k)=0,\ 0<r<\infty; \\ y_l(0;k)=0; \\ F_l(k;\hat{R})=0; \\ G_l(k;\hat{R})=0. \end{cases} \tag{1.18}$$

This is a two-way initial value problem starting at $r=\hat{R}$ inward and outward. The eigenvalue k passes through to the infinity by the uniqueness of the ODE and defines the far-field patterns near infinity. There is an one-to-one correspondence between the far-field patterns and the radiating solution of the Helmholtz equation. The $y_l(r;k)$ depends on the incident angle $\hat{x}$. Most important of all, we will examine the zero set of $y_l(0;k)=0$ which constitutes the eigenvalues of (1.18). The solutions $\{y_l(r;k)\}_{l\geq 0}$ is a family of entire functions of exponential type [5, 6, 7, 30]. For each $l\geq 0$, it behaves like sine functions in complex plane with zero set asymptotically approaching the zero set of sine functions for each incident direction. The Weyl's law of the eigenvalues of (1.18) in many settings are found in [10, 12, 13] as a direct consequence of the Cartwright-Levinson theory in value distribution theory [3, 8, 9, 23, 26, 27]. In particular, we can find that the density of the zero set for each incident direction is related to the radius $\hat{R}$ as a spectral invariant. Rellich's lemma indicates that all perturbations behave like spherical waves near the infinity, by which we prove **a special case** of Schiffer's conjecture.

Theorem 1.1. *Let D be a starlike domain as assumed in (1.1) under radiation condition (1.2). If there is an eigenvalue $k_0^2\in\mathbb{R}^+$, $k_0^2\geq 1$ of (1.1), then D is an open ball centered at the origin.*

2 Singular Sturm-Liouville Theory

Here we collect the asymptotic behaviors for $y_l(r;k)$ and $y_l'(r;k)$. For $l\geq 0$, we apply the results from [5, 6, 7, 30]. Let $z_l(\xi;k)$ be the solution of

$$\begin{cases} -z_l''(\xi)+\frac{l(l+1)z_l(\xi)}{\xi^2}+p(\xi)z_l(\xi)=k^2z_l(\xi); \\ z_l(1;k)=-b;\ z_l'(1;k)=a,\ a,\ b\in\mathbb{R}, \end{cases} \tag{2.1}$$

where $p(\xi)$ is square integrable; the real number $l\geq -1/2$. In general,

$$|z_l(\xi;k)+b\cos k(1-\xi)+a\frac{\sin k(1-\xi)}{k}|\leq\frac{K(\xi)}{|k|}\exp\{|\Im k|[1-\xi]\},\ |k|\geq 1, \tag{2.2}$$

where

$$K(\xi) \leq \exp\{\int_{\xi}^{1} \frac{|l(l+1)|}{t^2} + |p(t)|dt\},\ 0 \leq \xi \leq 1. \tag{2.3}$$

This explains the behaviors of solutions $z_l(\xi;k)$ and $z_l'(\xi;k)$ for all l in unit interval. For its application in (1.16) and (1.17), we take

$$b = -\hat{R};\ a = 1$$

for each incident direction, and the problem (2.1) in interval $[0, \hat{R}]$.

Outside the domain D, we consider (1.18) as an initial problem starting at $\hat{R}$ to the infinity. If $p(\xi) \equiv 0$, then we consider the following special case:

$$v_l''(\xi) + [k^2 - \frac{l(l+1)}{\xi^2}]v_l(\xi) = 0. \tag{2.4}$$

The solutions of (2.4) are essentially Bessel's functions with a basis of two elements. The variation of parameters formula leads to the following asymptotic expansions: For $\xi > 0$ and $\Re k \geq 0$, there is a constant C so that

$$|v_l(\xi,k) - \frac{\sin\{k\xi - l\frac{\pi}{2}\}}{k^{l+1}}| \leq C|k|^{-(l+1)}\frac{\exp\{|\Im k|\xi\}}{|k\xi|}; \tag{2.5}$$

$$|v_l'(\xi,k) - \frac{\cos\{k\xi - l\frac{\pi}{2}\}}{k^{l}}| \leq C|k|^{-l}\frac{\exp\{|\Im k|\xi\}}{|k\xi|}. \tag{2.6}$$

We refer these estimates to [6, Lemma 3.2, Lemma 3.3], and the we find that a solution of the initial value problem of (2.4) is a linear combination of (2.5) and (2.6).

3 Cartwright-Levinson Theory

We take the following vocabularies from entire function theory [3, 8, 9, 23, 26, 27] to describe the asymptotic behavior of the eigenvalues of (1.18).

Definition 3.1. Let $f(z)$ be an integral function of order ρ, $N(f,\alpha,\beta,r)$ be the number of the zeros of $f(z)$ inside the angle $[\alpha,\beta]$, and $|z| \leq r$. We define the density function as

$$\Delta_f(\alpha,\beta) := \lim_{r\to\infty} \frac{N(f,\alpha,\beta,r)}{r^\rho}, \tag{3.1}$$

and

$$\Delta_f(\beta) := \Delta_f(\alpha_0,\beta), \tag{3.2}$$

with some fixed $\alpha_0 \notin E$, in which E is at most a countable set [26, 27].

Let us define

$$\hat{\Delta}(\xi) := \Delta_{z_l(\xi;k)}(-\epsilon,\epsilon),\ b = -\hat{R}, \tag{3.3}$$

as the density of the zero set along $\hat{x}$.

Lemma 3.2. *The entire functions $y_l(\xi;k)$ and $y_l'(\xi;k)$ are of order one and of type $\hat{R} - \xi$.*

Proof. From (2.2), we have

$$y_l(\xi;k) = -\hat{R}\cos k(\hat{R}-\xi) - \frac{\sin k(\hat{R}-\xi)}{k} + O(\frac{K(\xi)}{|k|}\exp\{|\Im k|[\hat{R}-\xi]\}),\ |k| \geq 1. \tag{3.4}$$

To find the type of an entire function, we compute the following definition of Lindelöf's indicator function [26, 27]

Definition 3.3. Let $f(z)$ be an integral function of finite order ρ in the angle $[\theta_1,\theta_2]$. We call the following quantity as the indicator of the function $f(z)$.

$$h_f(\theta) := \lim_{r\to\infty} \frac{\ln|f(re^{i\theta})|}{r^\rho},\ \theta_1 \leq \theta \leq \theta_2. \tag{3.5}$$

We find that if $k = |k|e^{i\theta}$, then

$$h_{y_l(\xi;k)}(\theta) = |(\hat{R} - \xi)\sin\theta|,\ \theta \in [0, 2\pi],\ 0 < \xi < \hat{R}. \tag{3.6}$$

When referring more details to [10, 12, 13, 9, 26, 27], we find more examples in [9, p. 70]. The maximal value of $h_{y_l(\xi;k)}(\theta)$ gives the type of an entire function [26, p. 72], which is $(\hat{R} - \xi)$. A similar proof holds for $y_l'(\xi; k)$. □

More importantly, the indicator function (3.6) leads to the following Cartwright's theory [26, p. 251].

Lemma 3.4. *We have the following asymptotic behavior of the zero set of $y_l(\xi; k)$.*

$$\hat{\Delta}(\xi) = \frac{\hat{R} - \xi}{\pi}.$$

Proof. We observe in (3.4) that $|y_l(\xi; k)|$ is bounded on the real axis. Hence, it is in Cartwright's class. All of the properties in [26, p. 251] hold. □

Letting $\xi = 0$, we obtain the eigenvalue density of (1.18) in $\mathbb{C}$. Moreover, they are all real.

Lemma 3.5. *The eigenvalues k of (1.18) are all real.*

Proof. For $l = 0$, the result is classic [6, 30]. In our case, $y_l(\xi; k)$ is real for $k \in 0i + \mathbb{R}$. Furthermore, the asymptotic behavior of (3.4) proves the lemma, which is a special case of Bernstein's theorem in entire function theory [19, Theorem 1]. A step-by-step proof is provided in [13, Lemma 2.6]. □

4 Proof of Theorem 1.1

Proof. Let k_0^2 be an eigenvalue of (1.1), as assumed in Theorem 1.1. Particularly, from (1.5) we have

$$u(x; k_0) = \sum_{l=0}^{\infty} \sum_{m=-l}^{m=l} a_{l,m}(r; k_0) Y_l^m(\hat{x}); \tag{4.1}$$

$$u_{l,m}(x; k_0) = a_{l,m}(r; k_0) Y_l^m(\hat{x}),\ \hat{x} \in \mathbb{S}^2, \tag{4.2}$$

in which the coefficient $a_{l,m}(r; k_0)$ does not depend on the incident direction $\hat{x} \in \mathbb{S}^2$ for sufficiently large $|x| := r$: The functions in (4.1) solve the Helmholtz equation in $r \geq R_0$. As a result of the uniqueness of the ODE (1.18), the solutions $y_l(r; k_0)$ extend both outward to the infinity and inward to the origin for all $l \geq 0$. For the given eigenvalue k_0^2, the equation (1.18) holds for all incident directions $\hat{x} \in \mathbb{S}^2$ and for all $l \geq 0$.

The representation in (4.1) is unique in $\mathbb{R}^3$: If there is another eigenvalue k' of (1.18) from incident angle $x' \neq \hat{x} \in \mathbb{S}^2$ with the solution

$$u'_{l,m}(x; k') := a'_{l,m}(r; k') Y_l^m(\hat{x}),$$

then the analytic continuation of Helmholtz equation [17, p. 18] implies that

$$a'_{l,m}(r; k') = a_{l,m}(r; k_0). \tag{4.3}$$

With the uniqueness of the ODE (1.11), k_0 or k' satisfies (1.18) individually along its own incident direction inward to the origin. Therefore, Lemma 3.4 provides an eigenvalue density

$$\hat{\Delta}(0) = \frac{\hat{R}}{\pi},\ \hat{x} \in \mathbb{S}^2.$$

The ODE (1.18) holds for all $\xi \geq \hat{R}$ and $l \geq 0$. In particular, we apply the estimates (2.5) and (2.6):

$$|v_l(\xi, k) - \frac{\sin\{k(\xi - \hat{R}) - l\frac{\pi}{2}\}}{k^{l+1}}| \leq C|k|^{-(l+1)} \frac{\exp\{|\Im k|\xi\}}{|k\xi|};$$

$$|v_l'(\xi, k) - \frac{\cos\{k(\xi - \hat{R}) - l\frac{\pi}{2}\}}{k^{l}}| \leq C|k|^{-l} \frac{\exp\{|\Im k|\xi\}}{|k\xi|}.$$

Therefore, the initial value problem (2.1), with $p \equiv 0$, $b = \hat{R}$, and $a = 1$, provides the asymptotic behavior for the solution:

$$y_l(\xi;k) = \hat{R}\frac{\cos\{k(\xi-\hat{R}) - l\frac{\pi}{2}\}}{k^l} + O\{\frac{\exp\{|\Im k|\xi\}}{k^l(k\xi)}\}.$$

That is,

$$k^l y_l(\xi;k) = \hat{R}\cos\{k(\xi-\hat{R}) - l\frac{\pi}{2}\}[1+O\{\frac{1}{k\xi}\}],\ \hat{R} < \xi < \infty, \tag{4.4}$$

outside the zeros of $\cos\{k(\xi-\hat{R}) - l\frac{\pi}{2}\}$. This is classic in Sturm-Liouville theory [5, 6, 30].

The given eigenvalue k_0 satisfies (1.18), for all $l \geq 0$ and all $\hat{x} \in \mathbb{S}^2$, and (4.4). Therefore,

$$k_0^l y_l(\xi;k_0) = \hat{R}\cos\{k_0(\xi-\hat{R}) - l\frac{\pi}{2}\}[1+O\{\frac{1}{k_0\xi}\}],\ 0 < \xi < \infty. \tag{4.5}$$

We choose $l \uparrow \infty$ and so $\xi \uparrow \infty$ such that $\xi = \hat{R} + \frac{l\pi}{2k_0} > R_0$ for any large R_0. Thus, for large l,

$$k_0^l y_l(\xi;k_0) = \hat{R} + O(\frac{1}{\xi}),\ \xi = \hat{R} + \frac{l\pi}{2k_0},\ |k_0| \geq 1. \tag{4.6}$$

Using (1.10), (1.5) and the uniqueness of the Helmholtz equation, as shown in (4.3), the far-field patterns [17, (2.49)] are asymptotically the same periodic functions for each $\hat{x} \in \mathbb{S}^2$. In particular, the boundary defining function $\hat{R}$ is constant to $\hat{x} \in \mathbb{S}^2$, and Theorem 1.1 is thus proven. ☐

Compliance with Ethical Standards The author declares there is no conflicts of interest regarding the publication of this paper. The research does not involve any human participant and/or animals, and no further informed consent is required.

References

[1] M. Agranovsky, Integral geometry and spectral analysis, In "Topics in mathematical analysis", World Scientific Publishing Co, (281–320) 2008.

[2] T. Aktosun, D. Gintides and V. G. Papanicolaou, The uniqueness in the inverse problem for transmission eigenvalues for the spherically symmetric variable-speed wave equation, Inverse Problems, V. 27, 115004 (2011).

[3] R.P. Boas, Entire functions, Academic Press, New York, 1954.

[4] F. Cakoni, D. Colton and D. Gintides, The interior transmission eigenvalue problem, SIAM J. Math. Anal. 42, 2912–2921 (2010).

[5] R. Carlson, Inverse spectral theory for some singular Sturm-Liouville problems, Journal of Differential Equations, V.106, 121–140 (1993).

[6] R. Carlson, A Borg-Levinson theorem for Bessel operators, Pacific Journal of Mathematics, Vol. 177, No. 1, 1–26 (1997).

[7] R. Carlson, Inverse Sturm-Liouville problems with a singularity at zero, Inverse problems, 10, 851–864 (1994).

[8] M. L. Cartwright, On the directions of Borel of functions which are regular and of finite order in an angle, Proc. London Math. Soc. Ser. 2 Vol. 38, 503–541 (1933).

[9] M. L. Cartwright, Integral functions, Cambridge University Press, Cambridge, 1956.

[10] L. -H. Chen, An uniqueness result with some density theorems with interior transmission eigenvalues, Forthcoming in "Applicable Analysis".

[11] L. -H. Chen, An Inverse Uniqueness on the Index of Refraction with a Transition Region, Forthcoming in "Applicable Analysis".

[12] L. -H. Chen, A uniqueness theorem on the eigenvalues of spherically symmetric interior transmission problem in absorbing medium, Complex Var. Elliptic Equ. 60, no. 2, 145–167 (2015).

[13] L. -H. Chen, On the inverse spectral theory in a non-homogeneous interior transmission problem, Forthcoming in "Complex Variables and Elliptic Equations".

[14] D. Colton and P. Monk, The inverse scattering problem for time harmonic acoustic waves in a penetrable medium, Q. Jl. Mech. appl. Math. Vol. 40, 189–212 (1987).

[15] D. Colton and P. Monk, The inverse scattering problem for time-harmonic acoustic waves in an inhomogeneous medium, Q. Jl. Mech. appl. Math. Vol. 41, 97–125 (1988).

[16] D. Colton, L. Päivärinta and J. Sylvester, The interior transmission problem, Inverse problems and imaging, V. 1, no. 1, 13–28 (2007).

[17] D. Colton and R. Kress, Inverse acoustic and electromagnetic scattering theory, 2rd ed. Applied mathematical science, V. 93, Springer–Verlag, Berlin, 2013.

[18] D. Colton, A. Kirsch, and L. Päivärinta, Farfield patterns for acoustic waves in an inhomogeneous medium, SIAM J. Math. Anal. 20, 1472–1483 (1989).

[19] R. Duffin and A. C. Schaeffer, Some properties of functions of exponential type, Bull. Amer. Math. Soc, Providence, 44, 236–240 (1938).

[20] V. Isakov, Inverse problems for partial differential equations, Applied Mathematical Sciences, V. 127, Springer-Verlag, New York, 1998.

[21] A. Kirsch, The denseness of the far field patterns for the transmission problem, IMA J. Appl. Math, 37, no. 3, 213–225 (1986).

[22] A. Kirsch, On the existence of transmission eigenvalues, Inverse problems and imaging, 3, 155–172 (2009).

[23] P. Koosis, The logarithmic integral I, Cambridge University Press, New York, 1997.

[24] E. Lakshtanov and B. Vainberg, Bounds on positive interior transmission eigenvalues, Inverse Problems, V. 28, No. 10, 0266–5611 (2012).

[25] E. Lakshtanov and B. Vainberg, Weyl type bound on positive interior transmission eigenvalues, Comm. Partial Differential Equations, 39, no. 9, 1729–1740 (2014).

[26] B. Ja. Levin, Distribution of zeros of entire functions, revised edition, Translations of mathematical mongraphs, American mathematical society, Providence, 1972.

[27] B. Ja. Levin, Lectures on entire functions, Translation of mathematical monographs, V. 150, AMS, Providence, 1996.

[28] H. Y. Liu, Schiffer's conjecture, interior transmission eigenvalues and invisibility cloaking: singular problem vs. nonsingular problem, Geometric analysis and integral geometry, , Contemp. Math, 598, Amer. Math. Soc., Providence, RI, 147–154(2013).

[29] J. R. McLaughlin and P. L. Polyakov, On the uniqueness of a spherically symmetric speed of sound from transmission eigenvalues, Jour. Differential Equations, 107, 351–382 (1994).

[30] J. Pöschel and E. Trubowitz, Inverse spectral theory, Academic Press, Orlando, 1987.

[31] B. P. Rynne and B. D. Sleeman, The interior transmission problem and inverse scattering from inhomogeneous media, SIAM J. Math. Anal. 22, no.6, 1755–1762 (1991).